

A robust layer that delivers:

- Prolific Egg Numbers
- Persistency of Lay
- Excellent Livability
- Strong in Challenging Environments
- Outstanding Tinted Egg Quality





## Performance Summary

REARING PERIOD (TO 17 WEEKS):	
Livability	98%
Feed Consumed	5.78–6.15 kg
Body Weight at 17 Weeks	1.44–1.48 kg
	5
LAYING PERIOD (TO 90 WEEKS):	
Percent Peak	94–97%
Hen-Day Eggs to 60 Weeks	255–268
Hen-Day Eggs to 90 Weeks	417–440
Hen-Housed Eggs to 60 Weeks	251–265
Hen-Housed Eggs to 90 Weeks	406–428
Livability to 60 Weeks	97%
Livability to 90 Weeks	93%
Days to 50% Production (from hatch)	146 days
Average Egg Weight at 26 Weeks	56–59 g/egg
Average Egg Weight at 32 Weeks	60–62 g/egg
Average Egg Weight at 70 Weeks	64–66 g/egg
Total Egg Mass per Hen-Housed (18–90 weeks)	26.0 kg
Body Weight at 32 Weeks	1.84–1.93 kg
Body Weight at 70 Weeks	1.90–1.98 kg
Freedom from Egg Inclusions	Excellent
Shell Strength	Excellent
Shell Color at 38 Weeks	46
Shell Color at 56 Weeks	43
Shell Color at 70 Weeks	41
Haugh Units at 38 Weeks	93.0
Haugh Units at 56 Weeks	86.9
Haugh Units at 70 Weeks	83.0
Average Daily Feed Consumption (18–90 weeks)	103–109 g/day per bird
Feed Conversion Rate, kg Feed/kg Eggs (20–60 weeks)	1.95–2.06
Feed Conversion Rate, kg Feed/kg Eggs (20–90 weeks)	1.97–2.08
Feed Utilization, kg Egg/kg Feed (20-60 weeks)	0.49–0.50
Feed Utilization, kg Egg/kg Feed (20-90 weeks)	0.48–0.49
Feed Consumption per Dozen Eggs (20–60 weeks)	1.46–1.54 kg
Feed Consumption per Dozen Eggs (20–90 weeks)	1.51–1.60 kg
Skin Color	Yellow
Condition of Droppings	Dry



Hy-Line International PO Box 310 , Dallas Center, IA 50063 U.S.A. www.hyline.com



© 2018 Hy-Line International