

Hy-Line[®]

GRAY

COMMERCIAL LAYERS CONVENTIONAL SYSTEMS

Higher profitability due to:

- Superior Egg Production
- Excellent Livability
- Highest Tinted Egg Quality



Hy-Line[®]

Performance Summary

REARING PERIOD (TO 17 WEEKS):	
Livability	97.8%
Feed Consumed	5.69–6.08 kg
Body Weight at 17 Weeks	1336–1399 g
LAYING PERIOD (TO 100 WEEKS):	
Percent Peak	95.6–98.6%
Hen-Day Eggs to 60 Weeks	264.4–275.3
Hen-Day Eggs to 100 Weeks	496.0–517.6
Hen-Housed Eggs to 60 Weeks	261.3–272.1
Hen-Housed Eggs to 100 Weeks	481.3–502.2
Livability to 60 Weeks	97.6%
Livability to 80 Weeks	95.1%
Livability to 100 Weeks	91.6%
Days to 50% Production (from hatch)	143 days
Egg Weight at 26 Weeks	54.5–57.0 g/egg
Egg Weight at 32 Weeks	57.7–60.4 g/egg
Egg Weight at 70 Weeks	61.1–64.0 g/egg
Egg Weight at 100 Weeks	61.8–64.8 g/egg
Total Egg Mass per Hen-Housed (18–80 weeks)	24.3 kg
Total Egg Mass per Hen-Housed (18–100 weeks)	31.1 kg
Body Weight at 26 Weeks	1784–1864 g
Body Weight at 32 Weeks	1853–1930 g
Body Weight at 70 Weeks	1925–2013 g
Body Weight at 100 Weeks	1933–2028 g
Freedom From Egg Inclusions	Excellent
Shell Strength	Excellent
Shell Color Score at 38 Weeks	52
Shell Color Score at 56 Weeks	48
Shell Color Score at 72 Weeks	46
Haugh Units at 38 Weeks	95.0
Haugh Units at 56 Weeks	87.0
Haugh Units at 70 Weeks	83.0
Average Daily Feed Consumption (19–100 weeks)	108.8–113.3 g / day per bird
Feed Conversion Rate, kg Feed/kg Eggs (20–60 weeks)	1.95–2.03
Feed Conversion Rate, kg Feed/kg Eggs (20–100 weeks)	2.02–2.11
Feed Utilization, kg Egg/kg Feed (20–60 weeks)	0.49–0.51
Feed Utilization, kg Egg/kg Feed (20–100 weeks)	0.47–0.49
Feed Consumption per Dozen Eggs (20–60 weeks)	1.44–1.49 kg
Feed Consumption per Dozen Eggs (20–100 weeks)	1.49–1.55 kg
Skin Color	Yellow
Condition of Droppings	Dry

