



*Hy-Line*



**A robust layer that delivers:**

- **Prolific Egg Numbers**
- **Persistency of Lay for Long Single-Cycle Production**
- **Excellent Egg Shell Strength**
- **Strong Performance Under Challenging Environments or Low Density Feed Rations**
- **Adaptability to Alternative Production Systems**
- **Calm Disposition for Easy Management**

***Hy-Line***<sup>®</sup>

**W-80**

### Performance Summary

#### REARING PERIOD (TO 17 WEEKS):

Livability	97%
Feed Consumed	5.12–5.66 kg
Body Weight at 17 Weeks	1.17–1.25 kg

#### LAYING PERIOD (TO 100 WEEKS):

Percent Peak	95–97%
Hen-Day Eggs to 60 Weeks	262.2–269.0
Hen-Day Eggs to 100 Weeks	482.9–501.7
Hen-Housed Eggs to 60 Weeks	253.3–259.8
Hen-Housed Eggs to 100 Weeks	455.6–473.1
Livability to 60 Weeks	94.1%
Livability to 100 Weeks	89.3%
Days to 50% Production (from hatch)	141
Egg Weight at 26 Weeks	55.7 g / egg
Egg Weight at 32 Weeks	59.5 g / egg
Egg Weight at 70 Weeks	63.8 g / egg
Egg Weight at 100 Weeks	65.0 g / egg
Total Egg Mass per Hen-Housed (18–100 weeks)	28.7 kg
Body Weight at 26 Weeks	1.52–1.61 kg
Body Weight at 32 Weeks	1.59–1.71 kg
Body Weight at 70 Weeks	1.62–1.74 kg
Body Weight at 100 Weeks	1.66–1.76 kg
Freedom From Egg Inclusions	Excellent
Shell Strength	Excellent
Haugh Units at 38 Weeks	88.1
Haugh Units at 56 Weeks	85.6
Haugh Units at 70 Weeks	83.7
Haugh Units at 100 Weeks	80.0
Average Daily Feed Consumption (18–100 weeks)	101–108 g / day per bird
Feed Conversion Rate, kg Feed/kg Eggs (20–60 weeks)	1.88
Feed Conversion Rate, kg Feed/kg Eggs (20–100 weeks)	2.01
Feed Utilization, kg Egg/kg Feed (20–60 weeks)	0.53
Feed Utilization, kg Egg/kg Feed (20–100 weeks)	0.50
Feed Consumption per 10 Eggs (20–60 weeks)	1.10–1.15 kg
Feed Consumption per 10 Eggs (20–100 weeks)	1.19–1.23 kg
Condition of Droppings	Dry



Hy-Line International  
 PO Box 310 , Dallas Center, IA 50063 U.S.A.  
[www.hyline.com](http://www.hyline.com)

