

Hy-Line[®]

W-80

Conventional Systems



Performance Guide



Use of the Performance Guide

The genetic potential of Hy-Line W-80 Commercial can only be realized if good poultry husbandry practices and management are used. This management guide outlines successful flock management programmes for Hy-Line Variety W-80 Commercial based on field experience compiled by Hy-Line International and using an extensive commercial layer flock database of Hy-Line flocks from all parts of the world. Hy-Line International Management Guides are periodically updated as new performance data and/or nutrition information become available.

The information and suggestions contained in this management guide should be used for guidance and educational purposes only, recognising that local environmental and disease conditions may vary and a guide cannot cover all possible circumstances. While every attempt has been made to ensure that the information presented is accurate and reliable at the time of publication, Hy-Line International cannot accept responsibility for any errors, omissions or inaccuracies in such information or management suggestions. Further, Hy-Line International does not warrant or make any representations or guarantees regarding the use, validity, accuracy, or reliability of, or flock performance or productivity resulting from the use of, or otherwise respecting, such information or management suggestions. In no event shall Hy-Line International be liable for any special, indirect or consequential damages or special damages whatsoever arising out of or in connection with the use of the information or management suggestions contained in this management guide.

Always consult hyline.com for the latest performance, nutrition, and management information.



**Hy-Line W-80
Online Management Guide**

Table of Contents

Variety Standards

Summary of Performance Standards	3
Rearing Period Performance Table	4
Production Period Performance Table	5–6
Production Period Space Recommendations	7
Performance Graph	7
Egg Quality	8
Egg Size Distribution	8–9

Management

Brooding Period

Brooding Temperatures and Lighting	9
Development of the Organ Systems in Pullets	10
Body Score Chart	10

Transition Period

Transition Period from Rear to Peak Egg Production	11
--	----

Lighting

Light Programs	12
Intermittent Lighting Program for Chicks	12

Nutrition

Rearing Period

Nutritional Recommendations	13
-----------------------------	----

Production Period

Nutritional Recommendations (Economical Performance)	14
Dietary Nutrient Concentrations (Economical Performance)	15
Nutritional Recommendations (Optimal Performance)	16
Dietary Nutrient Concentrations (Optimal Performance)	17
Vitamins and Trace Minerals	18
Water Quality	19

Summary of Performance Standards

REARING PERIOD (TO 18 WEEKS):	
Livability	96.7%
Feed Consumed	6.03 kg
Body Weight at 17 Weeks	1218–1296 g
LAYING PERIOD (TO 100 WEEKS):	
Percent Peak	94.0–97.9%
Hen-Day Eggs to 60 Weeks	257.7–271.9
Hen-Day Eggs to 100 Weeks	490.5–514.4
Hen-Housed Eggs to 60 Weeks	249.6–263.4
Hen-Housed Eggs to 100 Weeks	463.6–486.4
Livability to 60 Weeks	94.7%
Livability to 100 Weeks	89.0%
Days to 50% Production (from hatch)	143
Egg Weight at 26 Weeks	53.1–57.9 g
Egg Weight at 32 Weeks	58.1–61.7 g
Egg Weight at 70 Weeks	61.8–65.6 g
Egg Weight at 100 Weeks	63.1–67.0 g
Total Egg Mass per Hen-Housed (19–100 weeks)	29.4 kg
Body Weight at 26 Weeks	1526–1623 g
Body Weight at 32 Weeks	1622–1726 g
Body Weight at 70 Weeks	1673–1780 g
Body Weight at 100 Weeks	1674–1781 g
Freedom From Egg Inclusions	Excellent
Shell Strength	Excellent
Haugh Units at 38 Weeks	88.1
Haugh Units at 56 Weeks	85.6
Haugh Units at 70 Weeks	83.7
Haugh Units at 100 Weeks	80.0
Average Daily Feed Consumption (19–100 weeks)	102.9–110.3 g/day per bird
Feed Conversion Rate, kg Feed/kg Eggs (20–60 weeks)	1.96–2.04
Feed Conversion Rate, kg Feed/kg Eggs (20–100 weeks)	2.04–2.12
Feed Utilization, kg Egg/kg Feed (20–60 weeks)	0.49–0.51
Feed Utilization, kg Egg/kg Feed (20–100 weeks)	0.47–0.49
Feed Consumption per Dozen Eggs (20–60 weeks)	1.37–1.42 kg
Feed Consumption per Dozen Eggs (20–100 weeks)	1.43–1.49 kg
Condition of Droppings	Dry

Rearing Period Performance Table

AGE (weeks)	MORTALITY Cumulative (%)	BODY WEIGHT (g)	WATER INTAKE (ml/bird/day)	FEED INTAKE (g/bird/day)	CUMULATIVE FEED INTAKE (g/bird to date)	UNIFORMITY %
1	1.00	68 – 72	18–30	12 – 15	84 – 105	>85%
2	1.32	122 – 130	24–40	16 – 20	196 – 245	
3	1.55	184 – 196	32–50	21 – 25	343 – 420	
4	1.65	255 – 271	39–60	26 – 30	525 – 630	>80%
5	1.80	331 – 352	47–70	31 – 35	742 – 875	
6	1.85	411 – 437	53–78	35 – 39	987 – 1,148	
7	1.95	518 – 551	57–88	38 – 44	1,253 – 1,456	>85%
8	2.05	605 – 644	63–94	42 – 47	1,547 – 1,785	
9	2.10	692 – 736	68–102	45 – 51	1,862 – 2,142	
10	2.25	776 – 826	72–108	48 – 54	2,198 – 2,520	
11	2.30	857 – 912	78–114	52 – 57	2,562 – 2,919	
12	2.40	934 – 994	83–120	55 – 60	2,947 – 3,339	
13	2.55	1003 – 1067	87–126	58 – 63	3,353 – 3,780	
14	2.70	1065 – 1133	92–132	61 – 66	3,780 – 4,242	
15	2.80	1120 – 1191	96–138	64 – 69	4,228 – 4,725	
16	2.90	1167 – 1241	102–146	68 – 73	4,704 – 5,236	
17	3.10	1218 – 1296	107–152	71 – 76	5,201 – 5,768	>90%
18	3.30	1252 – 1332	113–164	75 – 82	5,726 – 6,342	

Production Period Performance Table

AGE (weeks)	% HEN- DAY Current	HEN-DAY EGGS Cumulative	HEN- HOUSED EGGS Cumulative	MORT- ALITY Cumulative (%)	BODY WEIGHT (g)	WATER INTAKE (ml/bird/day)	FEED INTAKE (g/bird/day)	HH EGG MASS Cumulative (kg)	AVG. EGG WT. (g/egg)
19	6.6–12.1	0.5–0.8	0.5–0.8	0.0	1293–1376	127–188	78–85	0.02–0.04	40.9–44.8
20	27.6–47.7	2.4–4.2	2.4–4.2	0.2	1342–1428	135–197	81–91	0.11–0.19	43.5–47.7
21	61.1–82.0	6.7–9.9	6.6–9.9	0.5	1374–1462	142–205	87–95	0.31–0.46	45.8–50.1
22	79.0–89.7	12.2–16.2	12.1–16.1	0.6	1402–1492	148–215	89–98	0.59–0.77	47.7–52.2
23	87.0–90.6	18.3–22.5	18.2–22.4	0.8	1441–1533	154–222	90–100	0.90–1.10	49.3–54.0
24	89.7–93.4	24.6–29.1	24.4–28.9	0.9	1462–1555	159–228	92–102	1.23–1.44	50.7–55.5
25	91.3–95.1	31.0–35.7	30.7–35.5	1.1	1496–1591	162–230	95–104	1.57–1.80	51.9–56.8
26	92.6–96.4	37.4–42.5	37.1–42.2	1.2	1526–1623	163–231	96–105	1.93–2.17	53.1–57.9
27	93.3–97.2	44.0–49.3	43.6–48.9	1.4	1541–1639	164–232	97–107	2.29–2.55	54.3–58.8
28	93.8–97.7	50.5–56.1	50.0–55.6	1.5	1568–1668	164–233	98–108	2.66–2.94	55.3–59.6
29	94.0–97.9	57.1–63.0	56.5–62.3	1.7	1590–1692	164–233	99–108	3.04–3.33	56.2–60.3
30	94.0–97.9	63.7–69.8	63.0–69.1	1.8	1603–1705	164–233	100–109	3.42–3.73	57.0–60.8
31	93.9–97.8	70.3–76.7	69.4–75.8	2.0	1613–1716	164–233	101–110	3.80–4.13	57.8–61.3
32	93.8–97.8	76.8–83.5	75.8–82.5	2.1	1622–1726	164–234	102–110	4.19–4.53	58.1–61.7
33	93.8–97.7	83.4–90.4	82.3–89.2	2.3	1631–1735	164–233	102–110	4.58–4.93	58.5–62.1
34	93.7–97.6	90.0–97.2	88.7–95.8	2.4	1638–1743	164–233	103–111	4.96–5.33	58.8–62.4
35	93.6–97.5	96.5–104.0	95.0–102.5	2.6	1641–1746	163–233	104–111	5.35–5.74	59.1–62.7
36	93.6–97.5	103.1–110.9	101.4–109.1	2.7	1644–1749	163–233	104–111	5.74–6.14	59.3–62.9
37	93.5–97.4	109.6–117.7	107.8–115.7	2.9	1646–1751	163–233	105–112	6.13–6.55	59.5–63.2
38	93.5–97.4	116.2–124.5	114.1–122.4	3.0	1647–1752	163–232	105–112	6.52–6.96	59.6–63.3
39	93.4–97.3	122.7–131.3	120.5–129.0	3.1	1648–1753	163–232	105–112	6.91–7.36	59.8–63.5
40	93.3–97.2	129.2–138.1	126.8–135.5	3.3	1651–1756	163–232	105–112	7.30–7.77	59.9–63.6
41	93.3–97.2	135.8–144.9	133.1–142.1	3.4	1653–1759	163–232	105–112	7.69–8.18	60.1–63.8
42	93.3–97.2	142.3–151.7	139.4–148.7	3.5	1656–1762	163–232	105–112	8.08–8.58	60.2–63.9
43	93.2–97.1	148.8–158.5	145.7–155.2	3.6	1658–1764	163–232	105–112	8.48–8.99	60.3–64.0
44	93.2–97.1	155.3–165.3	152.0–161.8	3.7	1660–1766	163–232	105–112	8.87–9.40	60.4–64.1
45	93.1–97.0	161.9–172.1	158.2–168.3	3.8	1662–1768	163–232	105–112	9.26–9.81	60.4–64.2
46	92.8–96.6	168.4–178.9	164.5–174.8	4.0	1664–1770	163–232	105–112	9.65–10.21	60.5–64.3
47	92.7–96.5	174.8–185.6	170.7–181.3	4.1	1665–1771	163–232	105–112	10.03–10.62	60.6–64.3
48	92.4–96.3	181.3–192.4	176.9–187.7	4.2	1666–1772	163–232	105–112	10.42–11.02	60.7–64.4
49	92.3–96.1	187.8–199.1	183.1–194.2	4.3	1667–1773	163–232	105–112	10.81–11.42	60.7–64.5
50	91.9–95.8	194.2–205.8	189.2–200.6	4.4	1668–1774	163–232	105–112	11.19–11.82	60.8–64.5
51	91.8–95.6	200.6–212.5	195.3–207.0	4.5	1669–1775	163–232	105–112	11.58–12.22	60.8–64.6
52	91.6–95.4	207.0–219.2	201.5–213.3	4.6	1669–1776	163–232	105–112	11.96–12.62	60.9–64.7
53	91.3–95.1	213.4–225.8	207.6–219.7	4.7	1669–1776	163–232	105–112	12.35–13.02	61.0–64.7
54	91.1–94.9	219.8–232.5	213.6–226.0	4.8	1670–1777	163–232	105–112	12.73–13.42	61.0–64.8
55	90.8–94.6	226.2–239.1	219.7–232.3	4.9	1671–1778	163–232	105–112	13.11–13.82	61.1–64.8
56	90.6–94.4	232.5–245.7	225.7–238.6	5.0	1671–1778	163–232	105–112	13.49–14.21	61.1–64.9
57	90.3–94.0	238.8–252.3	231.7–244.8	5.1	1671–1778	163–232	105–112	13.87–14.61	61.2–64.9
58	90.0–93.8	245.1–258.8	237.7–251.0	5.1	1672–1779	163–232	105–112	14.24–15.00	61.2–65.0

Production Period Performance Table (cont.)


AGE (weeks)	% HEN-DAY Current	HEN-DAY EGGS Cumulative	HEN- HOUSED EGGS Cumulative	MORT- ALITY Cumulative (%)	BODY WEIGHT (g)	WATER INTAKE (ml/bird/day)	FEED INTAKE (g/bird/day)	HH EGG MASS Cumulative (kg)	AVG. EGG WT. (g/egg)
59	89.8–93.5	251.4–265.4	243.6 –257.3	5.2	1672–1779	163–232	105–112	14.62 – 15.39	61.3–65.0
60	89.5–93.3	257.7–271.9	249.6 –263.4	5.3	1672–1779	163–232	105–112	14.99 – 15.78	61.3–65.1
61	89.2–92.9	263.9–278.4	255.5 –269.6	5.4	1672–1779	163–232	105–112	15.37 – 16.17	61.3–65.1
62	88.9–92.6	270.1–284.9	261.3 –275.7	5.5	1673–1780	163–232	105–112	15.74 – 16.56	61.4–65.2
63	88.6–92.3	276.3–291.4	267.2 –281.8	5.6	1673–1780	163–232	105–112	16.11 – 16.95	61.4–65.2
64	88.4–92.0	282.5–297.8	273.0 –287.9	5.8	1673–1780	163–232	105–112	16.48 – 17.33	61.5–65.3
65	88.1–91.8	288.7–304.2	278.8 –293.9	5.9	1673–1780	163–232	105–112	16.85 – 17.71	61.5–65.3
66	87.8–91.5	294.8–310.6	284.6 –299.9	6.0	1673–1780	163–232	105–112	17.22 – 18.10	61.6–65.4
67	87.5–91.1	301.0–317.0	290.4 –305.9	6.2	1673–1780	163–232	105–112	17.58 – 18.48	61.6–65.4
68	87.2–90.8	307.1–323.4	296.1 –311.9	6.3	1673–1780	163–232	105–112	17.94 – 18.85	61.7–65.5
69	86.9–90.5	313.2–329.7	301.8 –317.8	6.5	1673–1780	163–232	105–112	18.31 – 19.23	61.7–65.5
70	86.6–90.2	319.2–336.0	307.4 –323.7	6.6	1673–1780	163–232	105–112	18.67 – 19.61	61.8–65.6
71	86.2–89.8	325.3–342.3	313.1 –329.6	6.7	1673–1780	163–232	105–112	19.03 – 19.98	61.8–65.6
72	86.0–89.6	331.3–348.6	318.7 –335.4	6.9	1674–1781	163–232	105–112	19.38 – 20.35	61.9–65.7
73	85.7–89.2	337.3–354.8	324.2 –341.2	7.0	1674–1781	163–232	105–112	19.74 – 20.72	61.9–65.7
74	85.3–88.9	343.2–361.0	329.8 –347.0	7.2	1674–1781	163–232	105–112	20.09 – 21.09	61.9–65.8
75	85.0–88.6	349.2–367.2	335.3 –352.7	7.3	1674–1781	163–232	105–112	20.44 – 21.46	62.0–65.8
76	84.7–88.2	355.1–373.4	340.8 –358.5	7.4	1674–1781	163–232	105–112	20.80 – 21.83	62.0–65.9
77	84.3–87.8	361.0–379.6	346.2 –364.1	7.6	1674–1781	163–232	105–112	21.14 – 22.19	62.1–65.9
78	84.1–87.6	366.9–385.7	351.7 –369.8	7.7	1674–1781	163–232	105–112	21.49 – 22.55	62.1–66.0
79	83.8–87.3	372.8–391.8	357.1 –375.4	7.9	1674–1781	163–232	105–112	21.84 – 22.91	62.2–66.0
80	83.4–86.9	378.6–397.9	362.4 –381.0	8.0	1674–1781	163–232	105–112	22.18 – 23.27	62.2–66.1
81	83.2–86.7	384.4–403.9	367.8 –386.6	8.2	1674–1781	163–232	105–112	22.53 – 23.63	62.3–66.1
82	82.9–86.3	390.2–410.0	373.1 –392.1	8.3	1674–1781	163–232	105–112	22.87 – 23.98	62.3–66.2
83	82.6–86.0	396.0–416.0	378.4 –397.7	8.4	1674–1781	163–232	105–112	23.21 – 24.34	62.3–66.2
84	82.2–85.6	401.8–422.0	383.7 –403.1	8.6	1674–1781	163–232	105–112	23.55 – 24.69	62.4–66.3
85	81.8–85.3	407.5–428.0	388.9 –408.6	8.7	1674–1781	163–232	105–112	23.88 – 25.04	62.4–66.3
86	81.5–84.9	413.2–433.9	394.1 –414.0	8.9	1674–1781	163–232	105–112	24.22 – 25.39	62.5–66.3
87	81.1–84.5	418.9–439.8	399.3 –419.4	9.0	1674–1781	163–232	105–112	24.55 – 25.74	62.5–66.4
88	80.8–84.2	424.5–445.7	404.4 –424.7	9.2	1674–1781	163–232	105–112	24.88 – 26.08	62.6–66.4
89	80.4–83.8	430.2–451.6	409.5 –430.0	9.3	1674–1781	163–232	105–112	25.21 – 26.43	62.6–66.5
90	80.0–83.4	435.8–457.4	414.6 –435.3	9.5	1674–1781	163–232	105–112	25.54 – 26.77	62.7–66.5
91	79.6–82.9	441.4–463.2	419.6 –440.6	9.6	1674–1781	163–232	105–112	25.87 – 27.11	62.7–66.6
92	79.3–82.6	446.9–469.0	424.6 –445.8	9.8	1674–1781	163–232	105–112	26.19 – 27.44	62.8–66.6
93	78.9–82.2	452.4–474.8	429.6 –451.0	9.9	1674–1781	163–232	105–112	26.51 – 27.78	62.8–66.7
94	78.6–81.9	457.9–480.5	434.5 –456.1	10.1	1674–1781	163–232	105–112	26.83 – 28.11	62.8–66.7
95	78.3–81.5	463.4–486.2	439.5 –461.3	10.2	1674–1781	163–232	105–112	27.15 – 28.45	62.9–66.8
96	77.9–81.2	468.9–491.9	444.4 –466.3	10.4	1674–1781	163–232	105–112	27.47 – 28.78	62.9–66.8
97	77.6–80.9	474.3–497.6	449.2 –471.4	10.5	1674–1781	163–232	105–112	27.78 – 29.11	63.0–66.9
98	77.3–80.6	479.7–503.2	454.1 –476.5	10.7	1674–1781	163–232	105–112	28.10 – 29.43	63.0–66.9
99	77.0–80.2	485.1–508.8	458.9 –481.5	10.9	1674–1781	163–232	105–112	28.41 – 29.76	63.1–67.0
100	76.7–79.9	490.5–514.4	463.6 –486.4	11.0	1674–1781	163–232	105–112	28.72 – 30.08	63.1–67.0

Production Period Space Recommendations

check local regulations concerning space requirements)

WEEKS OF AGE																									
31720304050607080																									
CONVENTIONAL AND COLONY CAGES																									
Floor Space																									
100–200 cm ² (50–100 birds / m ²)		310 cm ² (32 birds / m ²)														490 cm ² (20 birds / m ²) – 750 cm ² (13 birds / m ²)									
Nipple/Cup																									
1 / 12 birds		1 / 8 birds														1 / 12 birds or access to 2 drinkers									
Feeders																									
5 cm / bird		8 cm / bird														7–12 cm / bird									

Performance Graph



Egg Quality and Egg Size Distribution

E.U. Standards–Weekly*


AGE (weeks)	HAUGH UNITS	BREAKING STRENGTH
20	90.6	4490
22	90.3	4480
24	90.1	4470
26	89.8	4450
28	89.5	4430
30	89.2	4410
32	89.0	4390
34	88.7	4360
36	88.4	4340
38	88.1	4330
40	87.8	4320
42	87.6	4310
44	87.3	4300
46	87.0	4290
48	86.7	4280
50	86.5	4270
52	86.2	4260
54	85.9	4250
56	85.6	4240
58	85.4	4230
60	85.1	4220
62	84.8	4210
64	84.5	4200
66	84.3	4190
68	84.0	4170
70	83.7	4150
72	83.4	4130
74	83.2	4110
76	82.9	4090
78	82.6	4070
80	82.3	4050
82	82.1	4030
84	81.8	4010
86	81.5	4000
88	81.2	3980
90	81.0	3960
92	80.8	3950
94	80.6	3940
96	80.4	3930
98	80.2	3920
100	80.0	3910

AGE (weeks)	AVERAGE EGG WEIGHT (g)	WEEKLY % VERY LARGE Over 73 g	WEEKLY % LARGE 63–73 g	WEEKLY % MEDIUM 53–63 g	WEEKLY % SMALL 43–53 g
20	45.6	0.88	0.01	2.03	97.08
22	50.0	0.83	0.11	23.30	75.76
24	53.1	0.80	0.81	50.31	48.08
26	55.5	0.87	3.54	68.14	27.45
28	57.4	0.92	9.21	76.14	13.73
30	58.9	0.92	16.18	76.32	6.58
32	59.9	0.90	22.44	73.37	3.30
34	60.6	0.88	27.34	69.99	1.78
36	61.1	0.87	31.09	66.99	1.05
38	61.5	0.87	34.02	64.45	0.66
40	61.8	0.87	36.43	62.25	0.45
42	62.0	0.88	38.52	60.29	0.32
44	62.2	0.89	40.40	58.48	0.24
46	62.4	0.90	42.15	56.77	0.19
48	62.5	0.91	43.80	55.14	0.15
50	62.7	0.92	45.39	53.56	0.12
52	62.8	0.94	46.91	52.05	0.11
54	62.9	0.95	48.37	50.59	0.09
56	63.0	0.96	49.77	49.19	0.08
58	63.1	0.97	51.10	47.85	0.07
60	63.2	0.98	52.37	46.59	0.07
62	63.3	0.99	53.55	45.39	0.06
64	63.4	1.00	54.67	44.27	0.06
66	63.5	1.01	55.71	43.23	0.06
68	63.6	1.01	56.67	42.26	0.05
70	63.7	1.02	57.57	41.36	0.05
72	63.8	1.03	58.39	40.54	0.05
74	63.9	1.03	59.15	39.77	0.05
76	63.9	1.03	59.85	39.07	0.04
78	64.0	1.04	60.49	38.43	0.04
80	64.1	1.04	61.08	37.84	0.04
82	64.2	1.04	61.62	37.30	0.04
84	64.3	1.05	62.11	36.81	0.04
86	64.4	1.05	62.56	36.35	0.04
88	64.5	1.05	62.98	35.94	0.04
90	64.6	1.05	63.36	35.55	0.04
92	64.7	1.05	63.71	35.20	0.04
94	64.8	1.05	64.03	34.88	0.04
96	64.9	1.05	64.32	34.59	0.04
98	65.0	1.06	64.60	34.31	0.04
100	65.1	1.06	64.85	34.06	0.04

* Distribution of egg sizes based on weekly (not cumulative) average egg weights.

Egg Size Distribution (cont.)

E.U. Standards–Weekly*




* Distribution of egg sizes based on weekly (not cumulative) average egg weights.


Brooding Temperature and Lighting Recommendations

AGE	0–3 days	4–7 days	8–14 days	15–21 days	22–28 days	29–35 days	36–42 days
AIR TEMP. (CAGE)	33–36°C	30–32°C	28–30°C	26–28°C	23–26°C	21–23°C	21°C
AIR TEMP. (FLOOR)	35–36°C	33–35°C	31–33°C	29–31°C	26–27°C	23–25°C	21°C
LIGHT INTENSITY	30–50 lux	33–35°C	28–30°C	26–28°C	23–26°C	21–23°C	21°C
LIGHT HOURS	22 hours or Intermittent Program	30–50 lux	31–33°C	26–28°C	23–26°C	21–23°C	21°C
		21 hours or Intermittent Program	25 lux	29–31°C	23–26°C	21–23°C	21°C
			20 hours	25 lux	26–27°C	21–23°C	21°C
				18 hours	25 lux	23–25°C	21°C
					16.5 hours	5–10 lux	21°C
						15 hours	5–10 lux
							13.5 hours

Development of the Organ Systems in Pullets




Body Score Chart



Transition Period from Rear to Peak Egg Production

Frequently formulate to changing feed consumption during transition period until feed consumption is consistent.




Pre-Peak


- Pre-Peak diets are intended for flocks with low feed intake and are fed for a limited period from first egg to the beginning of peak production. The nutrient specification of the Pre-Peak diet should be dense enough to allow for lower feed intake and also cater to the increased nutritional needs of the bird entering egg production. Continue to feed the Pre-Peak until feed intake has developed sufficiently to allow transition to the Peak diet.
- If utilized until no more than 50–70% HD, a Pre-Peak diet with reduced energy concentration can be beneficial to stimulate feed intake. Pre-Peaking diets are useful in situations where local conditions may result in reduced feed intake, such as hot climates where feed intake may be depressed.
- Increasing the vitamins and trace mineral inclusion to 30% can be useful to cope with the lower feed intake during the Pre-Peak phase.

Light Programs

LIGHT-CONTROLLED HOUSING




OPEN-SIDED HOUSING




Intermittent Lighting Program for Chicks

- Preferred lighting technique.
- Use from 0–7 days (can be used up to 14 days of age).
- Intermittent dark periods provide rest periods for chicks.
- Synchronises chicks' activities and feedings.
- Establishes more natural behaviour of rest and activity.
- May improve 7-day livability and pullet body weight.
- Some dark periods may be shortened or removed to accommodate work schedules.



Rearing Period Nutritional Recommendations



¹ Body weights are approximate. Ages shown are a guide only. Please note that at time of transfer, there will be some loss in body weight (normally 10–12%) due to reduced water intake.

² Do not feed Pre-Lay Diet earlier than 15 weeks of age. Do not feed Pre-Lay later than first egg as it contains insufficient calcium to support egg production.

³ Recommended energy range is based on raw material energy values shown in feed ingredient table at back of this guide. It is important that target concentrations of dietary energy are adjusted according to energy system applied to raw material matrix.


⁴ Recommendation for Total Amino Acids is only appropriate to corn and soybean meal diet. Where diets utilise other ingredients, recommendations for Standardised Ileal Digestible Amino Acids must be followed.

⁵ Diets should always be formulated to provide required intake of amino acid. Concentration of crude protein in diet will vary with raw material used. Crude protein value provided is an estimated typical value only.

⁶ Calcium should be supplied as fine calcium carbonate (mean particle size less than 2 mm). Coarse limestone (2–4 mm) can be introduced in Pre-Lay Diet at up to 50% of total limestone.

⁷ Where other phosphorus systems are used, diets should contain recommended minimum level of available phosphorus.

Production Period Nutritional Recommendations for Economical Performance^{1,2}



Production Period Dietary Nutrient Concentrations for Economical Performance^{1,2}

FEEDING PHASE PRODUCTION	PEAKING ³ First egg until production drops 2% below peak					LAYER 2 2% below peak to 92%					LAYER 3 91–88%					LAYER 4 87–83%					LAYER 5 Less than 83%				
	RECOMMENDED CONCENTRATION																								
Metabolizable energy ⁴ , kcal/bird/day	295 / 310					290 / 305					285 / 300					280 / 295					275 / 290				
Metabolizable energy ⁴ , MJ/bird/day	1.23 / 1.30					1.21 / 1.28					1.19 / 1.26					1.17 / 1.23					1.15 / 1.21				
NUTRITION	FEED CONSUMPTION (*Typical Feed Consumption)																								
	g/day per bird					85	90	95*	100	105	95	100	105*	110	115	95	100	105*	110	115	95	100	105*	110	115
	Standardized Ileal Digestible Amino Acids																								
Lysine, %	0.96	0.91	0.86	0.82	0.78	0.90	0.85	0.81	0.77	0.74	0.83	0.79	0.75	0.72	0.69	0.80	0.76	0.72	0.69	0.66	0.77	0.73	0.70	0.66	0.63
Methionine, %	0.48	0.46	0.43	0.41	0.39	0.45	0.42	0.40	0.38	0.36	0.41	0.39	0.37	0.36	0.34	0.40	0.38	0.36	0.34	0.33	0.38	0.36	0.34	0.33	0.31
Methionine+Cystine, %	0.88	0.83	0.79	0.75	0.71	0.81	0.77	0.73	0.69	0.66	0.75	0.71	0.68	0.65	0.62	0.72	0.68	0.65	0.62	0.59	0.69	0.66	0.63	0.60	0.57
Threonine, %	0.68	0.64	0.60	0.57	0.55	0.63	0.60	0.57	0.54	0.52	0.58	0.55	0.53	0.50	0.48	0.56	0.53	0.51	0.48	0.46	0.54	0.51	0.49	0.46	0.44
Tryptophan, %	0.21	0.20	0.19	0.18	0.17	0.20	0.19	0.18	0.17	0.16	0.18	0.17	0.17	0.16	0.15	0.18	0.17	0.16	0.15	0.15	0.17	0.16	0.15	0.15	0.14
Arginine, %	1.00	0.95	0.90	0.85	0.81	0.94	0.89	0.84	0.80	0.77	0.87	0.82	0.78	0.75	0.71	0.83	0.79	0.75	0.72	0.69	0.80	0.76	0.72	0.69	0.66
Isoleucine, %	0.79	0.75	0.71	0.67	0.64	0.73	0.69	0.66	0.62	0.60	0.67	0.63	0.60	0.57	0.55	0.64	0.61	0.58	0.55	0.53	0.61	0.58	0.56	0.53	0.51
Valine, %	0.87	0.82	0.78	0.74	0.70	0.80	0.76	0.72	0.69	0.66	0.73	0.70	0.66	0.63	0.60	0.70	0.67	0.64	0.61	0.58	0.68	0.64	0.61	0.58	0.56
	Total Amino Acids ⁵																								
Lysine, %	1.06	1.00	0.95	0.90	0.86	0.99	0.93	0.89	0.84	0.81	0.91	0.87	0.82	0.79	0.75	0.88	0.83	0.79	0.76	0.72	0.84	0.80	0.76	0.73	0.69
Methionine, %	0.52	0.49	0.46	0.44	0.42	0.48	0.45	0.43	0.41	0.39	0.44	0.42	0.40	0.38	0.37	0.43	0.41	0.39	0.37	0.35	0.41	0.39	0.37	0.35	0.34
Methionine+Cystine, %	0.99	0.94	0.89	0.84	0.80	0.91	0.87	0.82	0.78	0.75	0.84	0.80	0.76	0.73	0.70	0.81	0.77	0.73	0.70	0.67	0.78	0.74	0.71	0.67	0.64
Threonine, %	0.79	0.75	0.71	0.68	0.64	0.74	0.70	0.67	0.64	0.61	0.69	0.65	0.62	0.59	0.57	0.66	0.63	0.60	0.57	0.54	0.63	0.60	0.57	0.55	0.52
Tryptophan, %	0.25	0.24	0.23	0.22	0.21	0.24	0.22	0.21	0.20	0.19	0.22	0.21	0.20	0.19	0.18	0.21	0.20	0.19	0.18	0.17	0.20	0.19	0.18	0.17	0.17
Arginine, %	1.08	1.02	0.97	0.92	0.87	1.01	0.95	0.91	0.86	0.82	0.93	0.88	0.84	0.80	0.77	0.89	0.85	0.81	0.77	0.74	0.86	0.82	0.78	0.74	0.71
Isoleucine, %	0.85	0.80	0.76	0.72	0.69	0.78	0.74	0.71	0.67	0.64	0.72	0.68	0.65	0.62	0.59	0.69	0.65	0.62	0.59	0.57	0.66	0.63	0.60	0.57	0.55
Valine, %	0.96	0.90	0.86	0.81	0.78	0.88	0.84	0.80	0.76	0.72	0.81	0.77	0.73	0.70	0.67	0.78	0.74	0.70	0.67	0.64	0.75	0.71	0.68	0.64	0.62
Crude protein ⁶ , %	20.71	19.56	18.53	17.60	16.76	19.33	18.32	17.40	16.57	15.82	17.79	16.90	16.10	15.36	14.70	17.16	16.30	15.52	14.82	14.17	16.53	15.70	14.95	14.27	13.65
Sodium, %	0.21	0.20	0.19	0.18	0.17	0.19	0.18	0.17	0.16	0.15	0.18	0.17	0.16	0.15	0.15	0.18	0.17	0.16	0.15	0.15	0.18	0.17	0.16	0.15	0.15
Chloride, %	0.21	0.20	0.19	0.18	0.17	0.19	0.18	0.17	0.16	0.15	0.18	0.17	0.16	0.15	0.15	0.18	0.17	0.16	0.15	0.15	0.18	0.17	0.16	0.15	0.15
Linoleic acid (C18:2 n-6), %	2.35	2.22	2.11	2.00	1.90	2.22	2.11	2.00	1.90	1.82	1.68	1.60	1.52	1.45	1.39	1.58	1.50	1.43	1.36	1.30	1.47	1.40	1.33	1.27	1.22
Choline, mg/kg	2118	2000	1895	1800	1714	2000	1895	1800	1714	1636	1895	1800	1714	1636	1565	1895	1800	1714	1636	1565	1895	1800	1714	1636	1565
CALCIUM AND PHOSPHORUS CHANGES BASED ON FEED INTAKE																									
	Weeks 18–32					Weeks 33–55					Weeks 56–72					Weeks 73–85					Weeks 86+				
Feed Consumption, g/day per bird	85	90	95	100	105	95	100	105	110	115	95	100	105	110	115	95	100	105	110	115	95	100	105	110	115
Calcium ^{7,8} , %	4.71	4.44	4.21	4.00	3.81	4.37	4.15	3.95	3.77	3.61	4.53	4.30	4.10	3.91	3.74	4.68	4.45	4.24	4.05	3.87	4.84	4.60	4.38	4.18	4.00
Phosphorus (available) ^{7,9} , %	0.53	0.50	0.47	0.45	0.43	0.44	0.42	0.40	0.38	0.37	0.42	0.39	0.38	0.36	0.34	0.39	0.37	0.35	0.34	0.32	0.36	0.34	0.33	0.31	0.30
Phosphorus (digestible), %	0.47	0.45	0.42	0.40	0.38	0.40	0.38	0.36	0.35	0.33	0.38	0.36	0.34	0.32	0.31	0.35	0.33	0.32	0.30	0.29	0.33	0.31	0.29	0.28	0.27

¹ All nutrient requirements are based on the [Feed Ingredient Tables](#).

² Crude protein, methionine+cystine, fat, linoleic acid, and / or energy may be changed to optimize egg size.

³ Peaking nutrient levels are calculated for birds at peak egg production. Prior to achieving peak egg production, the nutrient requirements will be lower.

⁴ A good approximation of the influence of temperature on energy needs is that for each 0.5°C change higher or lower than 22°C, subtract or add about 1.8 kcal /bird /day, respectively.

⁵ Recommendation for Total Amino Acids is only appropriate to corn and soybean meal diet. Where diets utilize other ingredients, recommendations for Standardized Ileal Digestible Amino Acids must be followed.


⁶ Diets should always be formulated to provide required intake of amino acid. Concentration of crude protein in diet will vary with raw material used. Crude protein value provided is an estimated typical value only.

⁷ Calcium and available phosphorus requirements are determined by flock age. When production remains higher and diets are fed for longer than ages shown, it is recommended to increase to calcium and phosphorus concentrations of next feeding phase.

⁸ Calcium carbonate particle size recommendation varies throughout lay. Refer to [Calcium Particle Size](#). Dietary calcium levels and fine to coarse ratio may need to be adjusted based on limestone solubility.

⁹ Where other phosphorus systems are used, diets should contain recommended minimum level of available phosphorus.

Production Period Nutritional Recommendations for Optimal Performance^{1,2}



FEEDING PHASE PRODUCTION	PEAKING ³ First egg until production drops 2% below peak					LAYER 2 2% below peak to 91%					LAYER 3 90–87%					LAYER 4 86–82%					LAYER 5 Less than 82%									
	RECOMMENDED CONCENTRATION																													
NUTRITION	Metabolizable energy ⁴ , kcal/bird/day					290 / 305					285 / 300					280 / 295					280 / 295					280 / 295				
	Metabolizable energy ⁴ , MJ/bird/day					1.21 / 1.28					1.19 / 1.26					1.17 / 1.23					1.17 / 1.23					1.17 / 1.23				
FEED CONSUMPTION (*Typical Feed Consumption)																														
g/day per bird	90	95	100*	105	110	100	105	110*	115	120	100	105	110*	115	120	100	105	110*	115	120	100	105	110*	115	120					
Standardized Ileal Digestible Amino Acids																														
Lysine, %	0.96	0.91	0.87	0.82	0.79	0.85	0.81	0.78	0.74	0.71	0.84	0.80	0.76	0.73	0.70	0.83	0.79	0.75	0.72	0.69	0.82	0.78	0.74	0.71	0.68					
Methionine, %	0.48	0.46	0.43	0.41	0.39	0.43	0.41	0.39	0.37	0.36	0.42	0.40	0.38	0.37	0.35	0.41	0.39	0.38	0.36	0.35	0.41	0.39	0.37	0.35	0.34					
Methionine+Cystine, %	0.88	0.84	0.80	0.76	0.72	0.78	0.74	0.71	0.67	0.65	0.76	0.72	0.69	0.66	0.63	0.75	0.71	0.68	0.65	0.62	0.73	0.70	0.67	0.64	0.61					
Threonine, %	0.67	0.64	0.61	0.58	0.55	0.60	0.57	0.54	0.52	0.50	0.59	0.56	0.53	0.51	0.49	0.58	0.55	0.53	0.50	0.48	0.57	0.54	0.52	0.50	0.48					
Tryptophan, %	0.21	0.20	0.19	0.18	0.17	0.19	0.18	0.17	0.16	0.16	0.19	0.18	0.17	0.16	0.15	0.18	0.17	0.17	0.16	0.15	0.18	0.17	0.16	0.16	0.15					
Arginine, %	1.00	0.95	0.90	0.86	0.82	0.89	0.84	0.81	0.77	0.74	0.87	0.83	0.79	0.76	0.73	0.86	0.82	0.78	0.75	0.72	0.85	0.81	0.77	0.74	0.71					
Isoleucine, %	0.79	0.75	0.71	0.68	0.64	0.69	0.66	0.63	0.60	0.58	0.67	0.64	0.61	0.58	0.56	0.66	0.63	0.60	0.58	0.55	0.65	0.62	0.59	0.57	0.54					
Valine, %	0.87	0.82	0.78	0.74	0.71	0.76	0.72	0.69	0.66	0.63	0.74	0.70	0.67	0.64	0.62	0.73	0.69	0.66	0.63	0.61	0.72	0.68	0.65	0.62	0.60					
Total Amino Acids ⁵																														
Lysine, %	1.05	1.00	0.95	0.90	0.86	0.93	0.89	0.85	0.81	0.78	0.92	0.88	0.84	0.80	0.77	0.91	0.86	0.82	0.79	0.76	0.89	0.85	0.81	0.78	0.74					
Methionine, %	0.52	0.49	0.47	0.44	0.42	0.46	0.44	0.42	0.40	0.38	0.45	0.43	0.41	0.39	0.38	0.45	0.42	0.40	0.39	0.37	0.44	0.42	0.40	0.38	0.37					
Methionine+Cystine, %	1.00	0.95	0.90	0.86	0.82	0.88	0.83	0.80	0.76	0.73	0.85	0.81	0.78	0.74	0.71	0.84	0.80	0.76	0.73	0.70	0.83	0.79	0.75	0.72	0.69					
Threonine, %	0.79	0.75	0.71	0.68	0.65	0.70	0.67	0.64	0.61	0.59	0.69	0.66	0.63	0.60	0.58	0.68	0.65	0.62	0.59	0.57	0.67	0.64	0.61	0.58	0.56					
Tryptophan, %	0.25	0.24	0.23	0.22	0.21	0.22	0.21	0.20	0.19	0.19	0.22	0.21	0.20	0.19	0.18	0.22	0.21	0.20	0.19	0.18	0.21	0.20	0.19	0.19	0.18					
Arginine, %	1.07	1.02	0.97	0.92	0.88	0.95	0.91	0.87	0.83	0.79	0.94	0.89	0.85	0.82	0.78	0.93	0.88	0.84	0.80	0.77	0.91	0.87	0.83	0.79	0.76					
Isoleucine, %	0.85	0.80	0.76	0.73	0.69	0.74	0.71	0.68	0.65	0.62	0.72	0.69	0.66	0.63	0.60	0.71	0.68	0.65	0.62	0.59	0.70	0.67	0.64	0.61	0.58					
Valine, %	0.95	0.90	0.86	0.82	0.78	0.84	0.80	0.76	0.73	0.70	0.82	0.78	0.74	0.71	0.68	0.80	0.76	0.73	0.70	0.67	0.79	0.75	0.72	0.69	0.66					
Crude protein ⁶ , %	20.44	19.37	18.40	17.52	16.73	18.18	17.31	16.52	15.80	15.15	17.95	17.10	16.32	15.61	14.96	17.73	16.88	16.11	15.41	14.77	17.50	16.67	15.91	15.22	14.58					
Sodium, %	0.21	0.20	0.19	0.18	0.17	0.18	0.17	0.16	0.16	0.15	0.18	0.17	0.16	0.16	0.15	0.18	0.17	0.16	0.16	0.15	0.18	0.17	0.16	0.16	0.15					
Chloride, %	0.21	0.20	0.19	0.18	0.17	0.18	0.17	0.16	0.16	0.15	0.18	0.17	0.16	0.16	0.15	0.18	0.17	0.16	0.16	0.15	0.18	0.17	0.16	0.16	0.15					
Linoleic acid (C18:2 n-6), %	2.22	2.11	2.00	1.90	1.82	2.00	1.90	1.82	1.74	1.67	1.60	1.52	1.45	1.39	1.33	1.50	1.43	1.36	1.30	1.25	1.40	1.33	1.27	1.22	1.17					
Choline, mg/kg	1778	1684	1600	1524	1455	1800	1714	1636	1565	1500	1800	1714	1636	1565	1500	1800	1714	1636	1565	1500	1800	1714	1636	1565	1500					

		CALCIUM AND PHOSPHORUS CHANGES BASED ON FEED INTAKE																								
		Weeks 18–32					Weeks 33–55					Weeks 56–72					Weeks 73–85					Weeks 86+				
		90	95	100	105	110	100	105	110	115	120	100	105	110	115	120	100	105	110	115	120	100	105	110	115	120
Feed Consumption, g/day per bird																										
Calcium ^{7,8} , %	4.44	4.21	4.00	3.81	3.64	4.15	3.95	3.77	3.61	3.46	4.30	4.10	3.91	3.74	3.58	4.45	4.24	4.05	3.87	3.71	4.84	4.60	4.38	4.18	3.83	
Phosphorus (available) ^{7,9} , %	0.50	0.47	0.45	0.43	0.41	0.42	0.40	0.38	0.37	0.35	0.38	0.38	0.36	0.34	0.33	0.37	0.35	0.34	0.32	0.31	0.34	0.33	0.31	0.30	0.29	
Phosphorus (digestible), %	0.45	0.42	0.40	0.38	0.36	0.38	0.36	0.35	0.33	0.32	0.36	0.34	0.32	0.31	0.30	0.33	0.32	0.30	0.29	0.28	0.31	0.29	0.28	0.27	0.26	

¹ All nutrient requirements are based on the [Feed Ingredient Tables](#).

² Crude protein, methionine+cystine, fat, linoleic acid, and / or energy may be changed to optimize egg size.

³ Peaking nutrient levels are calculated for birds at peak egg production. Prior to achieving peak egg production, the nutrient requirements will be lower.

⁴ A good approximation of the influence of temperature on energy needs is that for each 0.5°C change higher or lower than 22°C, subtract or add about 1.8 kcal /bird /day, respectively.

⁵ Recommendation for Total Amino Acids is only appropriate to corn and soybean meal diet. Where diets utilize other ingredients, recommendations for Standardized Ileal Digestible Amino Acids must be followed.

⁶ Diets should always be formulated to provide required intake of amino acid. Concentration of crude protein in diet will vary with raw material used. Crude protein value provided is an estimated typical value only.

⁷ Calcium and available phosphorus requirements are determined by flock age. When production remains higher and diets are fed for longer than ages shown, it is recommended to increase to calcium and phosphorus concentrations of next feeding phase.

⁸ Calcium carbonate particle size recommendation varies throughout lay. Refer to [Calcium Particle Size](#). Dietary calcium levels and fine to coarse ratio may need to be adjusted based on limestone solubility.

⁹ Where other phosphorus systems are used, diets should contain recommended minimum level of available phosphorus.

Vitamins and Trace Minerals

ITEM ^{1,2,3,4}	IN 1000 KG COMPLETE DIET	
	Rearing Period	Production Period
Vitamin A, IU	10,000,000	8,000,000
Vitamin D ₃ ⁵ , IU	3,300,000	3,300,000
Vitamin E, g	30.00	25.00
Vitamin K (menadione), g	3.50	3.00
Thiamin (B ₁), g	2.20	2.50
Riboflavin (B ₂), g	6.60	5.50
Niacin (B ₃) ⁶ , g	40.00	30.00
Pantothenic acid (B ₅), g	10.00	10.00
Pyridoxine (B ₆), g	4.50	5.00
Biotin (B ₇), mg	100.00	75.00
Folic acid (B ₉), g	1.00	0.90
Cobalamine (B ₁₂), mg	23.00	23.00
Manganese ⁷ , g	100.00	100.00
Zinc ⁷ , g	85.00	80.00
Iron ⁷ , g	30.00	40.00
Copper ⁷ , g	15.00	8.00
Magnesium ⁷ , g	600.00	500.00
Iodine, g	1.50	1.20
Selenium ⁷ , g	0.25	0.25

¹ Minimum recommendations for rearing and laying periods. Local regulations may limit dietary content of individual vitamins or minerals. Levels of 150-200mg/kg of Vitamin C can be beneficial during periods of stress.

² Store premixes according to supplier's recommendations and observe 'use by' dates to ensure vitamin activity is maintained. Inclusion of antioxidant may improve premix stability.

³ Vitamin and mineral recommendations vary according to activity.

⁴ Where heat treatment is applied to diet, higher levels of vitamins may be required. Consult with vitamin supplier regarding stability through individual production processes.

⁵ A proportion of Vitamin D₃ can be supplemented as 25-hydroxy D₃ according to supplier's recommendations and applicable limits.

⁶ Higher levels of Niacin are recommended in non-cage systems.

⁷ Greater bioavailability and productivity may be possible with use of chelated mineral sources.

Drinking Water Quality for Poultry

ITEM	MAXIMUM CONCENTRATION (ppm or mg/L)*	
Nitrate NO_3^- ¹	25	Older birds will tolerate higher levels up to 20 ppm. Stressed or disease challenged birds may be more sensitive to effects of Nitrate.
Nitrate Nitrogen ($\text{NO}_3\text{-N}$) ¹	6	
Nitrite NO_2^- ¹	4	Nitrite is considerably more toxic than Nitrate, especially for young birds, where 1 ppm Nitrite may be considered toxic.
Nitrite Nitrogen ($\text{NO}_2\text{-N}$) ¹	1	
Total dissolved solids ²	1000	Levels up to 3000 ppm may not affect performance but could increase manure moisture.
Chloride (Cl^-) ¹	250	Levels as low as 14 mg may be problematic if sodium is higher than 50 ppm.
Sulphate (SO_4^-) ¹	250	Higher levels may be laxative.
Iron (Fe) ¹	<0.3	Higher levels result in bad odour and taste.
Magnesium (Mg) ¹	125	Higher levels may be laxative. Levels above 50 ppm may be problematic if sulphate levels are high.
Potassium (K) ²	20	Higher levels may be acceptable depending on sodium level, alkalinity, and pH.
Sodium (Na) ^{1,2}	50	Higher concentration is acceptable but concentrations above 50 ppm should be avoided if high levels of chloride, sulphate, or potassium exist.
Manganese (Mn) ³	0.05	Higher levels may be laxative.
Arsenic (As) ²	0.5	
Fluoride (F^-) ²	2	
Aluminium (Al) ²	5	
Boron (B) ²	5	
Cadmium (Cd) ²	0.02	
Cobalt (Co) ²	1	
Copper (Cu) ¹	0.6	Higher levels result in bitter taste.
Lead (Pb) ¹	0.02	Higher levels are toxic.
Mercury (Hg) ²	0.003	Higher levels are toxic.
Zinc (Zn) ¹	1.5	Higher levels are toxic.
pH ¹	5–7	Birds may adapt to lower pH. Below pH 5 may reduce water intake and corrode metal fittings. Above pH 8 may reduce intake and reduce effectiveness of water sanitation.
Total bacteria counts ³	1000 CFU/ml	This is likely to indicate dirty water.
Total Coliform bacteria ³	50 CFU/ml	
Faecal Coliform bacteria ³	0 CFU/ml	
Oxygen Reduction Potential (ORP) ³	650–750 mEq	The ORP range at which 2–4 ppm of free chlorine will effectively sanitise water at a favourable pH range of 5–7.

* Limits may be lower as interactions exist between magnesium and sulphate; and between sodium, potassium, chloride, and sulphate.

¹ Carter & Sneed, 1996. Drinking Water Quality for Poultry, Poultry Science and Technology Guide, North Carolina State University Poultry Extension Service. Guide no. 42

² Marx and Jaikaran, 2007. Water Analysis Interpretation. Agri-Facts, Alberta Ag-Info Centre. Refer to <http://www.agric.gov.ab.ca/app84/rwqit> for online Water Analysis Tool

³ Watkins, 2008. Water: Identifying and Correcting Challenges. Avian Advice 10(3): 10–15 University of Arkansas Cooperative Extension Service, Fayetteville

Always consult
[hyline.com](https://www.hyline.com) for the
latest performance,
nutrition, and
management
information.



Hy-Line W-80 Online Management Guide

RESOURCES AVAILABLE AT [WWW.HYLINE.COM](https://www.hyline.com)

[Corporate Information](#) | [Technical Updates](#) | [Videos](#) | [Interactive Management Guides](#)
[Hy-Line International Lighting Program](#) | [Hy-Line EggCel](#) | [Body Weight Uniformity Calculator](#)

TECHNICAL UPDATES

Diseases

An Overview of Focal Duodenal Necrosis (FDN)

MG Control in Commercial Layers

Colibacillosis in Layers: An Overview

Fowl Pox in Layers

Avian Urolithiasis (Visceral Gout)

Infectious Bursal Disease (IBD, Gumboro)

Fatty Liver Hemorrhagic Syndrome

Infectious Laryngotracheitis (ILT)

Intestinal Dilation Syndrome (IDS)

Newcastle Disease

Mycoplasma Synoviae (MS)

Low Pathogenic Avian Influenza (LPAI)

Diagnostic Samples and Breeder Flock Monitoring

Salmonella, *Mycoplasma*, and Avian Influenza
Monitoring in Parent Breeder Flocks

Proper Collection and Handling of Diagnostic Samples

Management

Growing Management of Commercial Pullets

Understanding the Role of the Skeleton in Egg Production

The Science of Egg Quality

Understanding Poultry Lighting

Understanding Heat Stress in Layers

Infrared Beak Treatment

Feed Granulometry and the Importance of
Feed Particle Size in Layers

Impact of Tarp Colour on Poultry Lighting

SPIDES (Short Period Incubation During Egg Storage)

Fly Management: Surveillance and Control

Optimising Egg Size in Commercial Layers

Vaccination Recommendations

Egg Drop Syndrome (EDS)

Managing Fully Beaked Flocks

Thiamin Deficiency in Pullets

Understanding Nesting Behaviour

Hy-Line International | www.hyline.com

Hy-Line is a brand name. ® Registered trademark of Hy-Line International.

© Copyright 2023 Hy-Line International.

80 STD ENG 061825

