

Hy-Line[®]

W-80

COMMERCIAL LAYERS ALTERNATIVE SYSTEMS

A robust layer that delivers:

- Docile Temperament for Alternative Systems
- Prolific Egg Numbers, Large or Small
- Persistency of Lay for Extended Laying Cycles
- Excellent Egg Shell Strength
- Excellent Nesting Behaviour



Hy-Line[®]

Performance Summary

REARING PERIOD (TO 17 WEEKS):	
Livability	96.20%
Feed Consumed	5.78–5.99 kg
Body Weight at 17 Weeks	1212–1269 g
LAYING PERIOD (TO 100 WEEKS):	
Percent Peak	93.9–97.7%
Hen-Day Eggs to 60 Weeks Hen-Day Eggs to 100 Weeks	258.1–268.7 494.8–515.0
Hen-Housed Eggs to 60 Weeks Hen-Housed Eggs to 100 Weeks	248.8–258.9 465.3–484.3
Livability to 60 Weeks Livability to 80 Weeks Livability to 100 Weeks	95.08% 92.85% 90.01%
Days to 50% Production (from hatch)	142
Egg Weight at 26 Weeks Egg Weight at 32 Weeks Egg Weight at 70 Weeks Egg Weight at 100 Weeks	54.0–55.7 g 58.5–60.3 g 62.5–64.4 g 62.3–64.2 g
Total Egg Mass per Hen-Housed (19–80 weeks) Total Egg Mass per Hen-Housed (19–100 weeks)	23.2–23.9 kg 29.6–30.5 kg
Body Weight at 26 Weeks Body Weight at 32 Weeks Body Weight at 70 Weeks Body Weight at 100 Weeks	1495–1570 g 1573–1657 g 1641–1756 g 1675–1814 g
Freedom From Egg Inclusions	Excellent
Shell Strength	Excellent
Haugh Units at 38 Weeks Haugh Units at 56 Weeks Haugh Units at 70 Weeks Haugh Units at 100 Weeks	88.1 85.6 83.7 80.0
Average Daily Feed Consumption (19–100 weeks)	109.4–113.3 g/day per bird
Feed Conversion Rate, kg Feed/kg Eggs (20–60 weeks) Feed Conversion Rate, kg Feed/kg Eggs (20–100 weeks)	1.99–2.07 2.10–2.19
Feed Utilization, kg Egg/kg Feed (20–60 weeks) Feed Utilization, kg Egg/kg Feed (20–100 weeks)	0.48–0.50 0.46–0.48
Feed Consumption per Dozen Eggs (20–60 weeks) Feed Consumption per Dozen Eggs (20–100 weeks)	1.46–1.52 kg 1.50–1.56 kg
Condition of Droppings	Dry

Hy-Line®

W-80



Hy-Line®

Hy-Line International | www.hyline.com

Hy-Line is a brand name. ® Registered trademark of
Hy-Line International. © Copyright 2026 Hy-Line International.